

MboI

5'-G A T C-3'
3'-C T A G-5'

| Cat. No. | Size |
|----------|-------------|
| E2283-01 | 200 units |
| E2283-02 | 1 000 units |

Reaction Temperature: 37°C

Inactivation Temperature (20 min): 65°C

Prototype: MboI

Source: *Moraxella bovis*

Package Contents:

- **MboI**
- **10x Reaction Buffer Medium**
- **BSA [100x]**
Added as separate component to prevent reaction buffer precipitation.
- **Dilution Buffer # 1**
Added for enzymes exceeding 10 U/μl in concentration. High protein concentration warrants optimal stability during prolonged storage. Use dilution buffer to prepare short term working stocks (5-10 U/μl, non-freezing at -20°C).

Storage Conditions: Store at -20°C

Double Digestion – Buffer Compatibility:

| Buffer | % Rel. Activity |
|--------|-----------------|
| Low | 75 |
| Medium | 100 |
| High | 100 |
| Acet | 100 |

Recommended Buffer: Medium
(or compatible third party buffers)

Restriction Enzyme Buffer Compatibility:
Both, enzyme and buffers are fully compatible to restrictases and buffer systems from other manufacturers and can be used along in double digestions. To obtain best results, consult the corresponding manuals of all involved products.

DNA Methylation:

No inhibition: dcm, EcoKI
Potential inhibition: CpG
Inhibition (Blocked): dam

Standard Reaction Protocol:

Mix the following reaction components:

- 1-2 μg pure DNA or 10 μl PCR product (≈0.1-2 μg DNA)
- 5 μl 10x Buffer Medium
- 0.5 μl BSA [100x]
- 1-2 U MboI (use 1 U / μg DNA, < 10 % React. Volume!)

Tips: Add enzyme as last component. Mix components well before adding enzyme. After enzyme addition, mix gently by pipetting. Do not vortex. High (excess) amounts of enzyme can greatly speed up the reaction.

@ 50 μl H₂O, nuclease free

Incubate for 1 h at 37°C

To obtain complete digestion of high molecular weight DNA, (e.g. plant genomic DNA), add excess amounts of enzyme and prolong the incubation time.

Stop reaction by alternatively

- (a) Addition of 2.1 μl EDTA pH 8.0 [0.5 M], final 20 mM or
- (b) Heat Inactivation
20 min at 65°C or
- (c) Spin Column DNA Purification
(e.g. EURx PCR/DNA CleanUp Kit, Cat.No. E3520) or
- (d) Gel Electrophoresis and Single Band Excision
(e.g. EURx AgaroseOut DNA Kit, Cat.No. E3540) or
- (e) Phenol-Chloroform Extraction or Ethanol Precipitation.

Non-optimal buffer conditions:

To compensate for the lack of enzyme activity, increase the amount of enzyme and / or reaction time accordingly. The following values may serve as orientation:

1. *Enzyme amount:* Instead of 1 U enzyme, use ~4 U of enzyme in buffers providing 25 % rel. activity, ~2 U in 50 %, ~1.5 U in 75 % or ~1 U in 100 %, respectively.
2. *Reaction time:* Increase by ~1.3-fold (75 % rel. activity), ~2-fold (50 %) or ~4-fold (25 %).

Unit Definition:

One unit is the amount of enzyme required to completely digest 1 μg of unmethylated Lambda DNA in 1 hr in a total reaction volume of 50 μl. Enzyme activity was determined in the recommended reaction buffer.

Reaction Buffer:

1 x Medium Buffer: 10 mM Tris-HCl (pH 7.5 at 37°C), 10 mM MgCl₂, 50 mM NaCl, 1 mM dithiothreitol.
To be supplemented with 100 μg/ml bovine serum albumin.

Note: blocked by *Dam* methylation

Storage Buffer:

10 mM Tris-HCl (pH 7.5 at 22°C), 50 mM NaCl, 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml bovine serum albumin, 0.01 % Triton™X-100 and 50 % (v/v) glycerol.

Quality Control:

All preparations are assayed for contaminating endonuclease, 3'-exonuclease, 5'-exonuclease/5'-phosphatase, as well as nonspecific single- and double-stranded DNase activities.