



HOT FIREPol[®] DNA Polymerase

(HOT START DNA Polymerase)

Cat. No.	Pack Size	Conc.
01-02-0000S	100 U SAMPLE	5 U/µl
01-02-00500	500 U	5 U/µl
01-02-01000	1000 U	5 U/µl

For in vitro use only.

Description:

HOT FIREPol[®] DNA Polymerase is a chemically modified FIREPol[®] DNA Polymerase. At ambient temperatures it is inactive, having no polymerization activity. HOT FIREPol[®] DNA Polymerase is activated by a 15 min incubation step at 95°C. This prevents extension of non-specifically annealed primers and primer-dimers formed at low temperatures during PCR setup. The enzyme has 5'-3' polymerization-dependent exonuclease replacement activity but lacks 3'-5' exonuclease activity.

Source:

Purified from an *E.coli* strain that carries an overproducing plasmid containing a modified gene of *Thermus aquaticus* DNA Polymerase.

Applications:

- Hot Start PCR
- DHPLC
- TA cloning

Reagents Provided:

- HOT FIREPol[®] DNA Polymerase
- HOT FIREPol[®] 10x Buffer B1 (Mg²⁺ and detergent free) Tris-HCl and (NH4)₂SO₄
- free) Tris-HCl and (NH₄)₂SO₄
 HOT FIREPol[®] 10x Buffer B2 (Mg²⁺ free) Tris-HCl, (NH₄)₂SO₄ and detergent
- **10x Solution S** Additive that facilitates amplification of difficult templates (e.g. GC-rich DNA templates). This solution should be used at a defined working concentration (1x, 2x or 3x solution).

Solution S is NOT a reaction buffer and should be used ONLY IF non-specific amplifications occur.

Concentration:

5 U/µl

Unit definition:

One unit is defined as the amount of the enzyme required to catalyze the incorporation of 10 nmol of dNTPs into an acid-insoluble form in 30 minutes at 74°C.

Storage and Dilution buffer:

50% glycerol (v/v), 20 mM Tris-HCl pH 8.7 at 25°C, 100 mM KCl, 0.1 mM EDTA and stabilizers.

Quality control:

The enzyme is free of nicking and priming activities, exonucleases and non-specific endonucleases. SDS/PAGE - 95 kD band, >98% pure. Activity and stability tested via thermo cycling. The error rate per nucleotide per cycle is ~ 2.5×10^{-5} ; the accuracy is ~ 4×10^{4} . Estimated half life at 95°C is 1.5 hours.

Shipping and Storage conditions:

Routine storage: -20°C

Shipping and temporary storage for up to 1 month at room temperature has no detrimental effects on the quality of HOT FIREPol[®] DNA Polymerase.

Recommended PCR Buffers:

Our standard buffers are:

- HOT FIREPol[®] 10x Buffer B1 (Mg²⁺ and detergent free) Tris-HCl and (NH₄)₂SO₄
- HOT FIREPol[®] 10x Buffer B2 (Mg²⁺ free) Tris-HCl, (NH₄)₂SO₄ and detergent

For further optimization please try our whole set of buffers available upon request:

- 10x Reaction buffer B (Mg²⁺ free) 0.8 M Tris-HCl, 0.2 M (NH₄)₂SO₄, 0.2% w/v Tween-20
- 10x Reaction buffer BD (Mg²⁺ and detergent free) 0.8 M Tris-HCl, 0.2 M (NH₄)₂SO₄

Recommended PCR reaction mix:

Component	Volume	Final conc.
HOT FIREPol [®] (5 U/µI)	0.4-1.0 µl	0.02-0.05 U/µl (2-5 U)
10x Buffer B1/B2,B/BD	10 µl	1x
25 mM MgCl ₂	6-10 µl	1.5-2.5 mM
20 mM dNTP mix	1 µl	200 µM
Primer Forward (10 pmol/µl)	1-3 µl	0.1-0.3 µM
Primer Reverse (10 pmol/µl)	1-3 µl	0.1-0.3 µM
DNA template	5-20 µl	5-100 ng/µl
10x Solution S Not for standard PCR	0, 10, 20 or 30 µl	1x, 2x or 3x
H ₂ O PCR grade	Up to 100 µl	
Total	100 µl	

REŞİT GALİP CADDESİ 74-7 06700 ÇANKAYA, ANKARA, TÜRKİYE T +90 312 447 22 79 / 80 F +90 312 447 22 07 W www.bmlabosis.com



Recommended PCR cycles:

Cycle step	Temp.	Time	Cycles
Initial denaturation	95°C	12-15 min	1
Denaturation	95°C	30-60 s	
Annealing	50-68°C	30-60 s	26-35
Elongation	72°C	1-4 min	
Final elongation	72°C	5-10 min	1

IMPORTANT: To activate the polymerase, include an incubation step at 95°C for 12 - 15 minutes at the beginning of the PCR cycle. Annealing temperature should be 2-6°C lower than the primer melting temperature. Elongation time should be ~1 min/1 kb.

Safety warnings and precautions:

This product and its components should be handled only by persons trained in laboratory techniques. It is advisable to wear suitable protective clothing, such as laboratory overalls, gloves and safety glasses. Care should be taken to avoid contact with skin or eyes. In case of contact with skin or eyes, wash immediately with water.

Some applications this product is used in may require a license which is not provided by the purchase of this product. Users should obtain the license if required.

Related products:

Product name	Pack size	Cat. No.
FIREPol [®] DNA Polymerase	500 U	01-01-00500
FIREPol [®] DNA Polymerase	1000 U	01-01-01000
FIREPol [®] DNA Polymerase	2000 U	01-01-02000
5x FIREPol[®] Master Mix (1.5 mM MgCl ₂ final conc.)	250 reactions	04-11-00115
5x FIREPol [®] Master Mix (2.5 mM MgCl ₂ final conc.)	250 reactions	04-11-00125
5x FIREPol [®] Master Mix Ready to Load (1.5 mM MgCl₂ final conc.)	250 reactions	04-12-00115
5x FIREPol [®] Master Mix Ready to Load (2.5 mM MgCl₂ final conc.)	250 reactions	04-12-00125
dNTP MIX (20 mM of each)	20 µmol	02-31-00020
dNTP MIX (20 mM of each)	100 µmol	02-31-00100
dNTP SET (100 mM)	4 x 25 µmol	02-21-00100
dNTP SET (100 mM)	4 x 100 µmol	02-21-00400